

## First record of the swimming sea cucumber *Enypniastes eximia* Théel, 1882 (Echinodermata: Holothuroidea) in Peruvian waters

### Primer registro del pepino de mar nadador *Enypniastes eximia* Theel, 1882 (Echinodermata: Holothuroidea) en aguas peruanas

Francisco Alonso Solís-Marín<sup>1</sup>, Yuri Hooker<sup>2</sup> and Alfredo Laguarda-Figuera<sup>1</sup>

<sup>1</sup> Laboratorio de Sistemática y Ecología de Equinodermos, Instituto de Ciencias del Mar y Limnología (ICML), Universidad Nacional Autónoma de México (UNAM). Apdo. Post. 70-305, México, D. F. 04510, México.

E-mail Francisco A. Solís-Marín: fasilis@cmarl.unam.mx

<sup>2</sup> Laboratorio de Biología Marina, Departamento de Ciencias Biológicas y Fisiológicas, Universidad Peruana Cayetano Heredia. Av. Honorio Delgado 430, Urb. Ingeniería, S.M.P. Lima, Perú.

#### Abstract

*Enypniastes eximia* Théel, 1882 (Echinodermata: Holothuroidea) a swimming sea cucumber is reported for the first time from Peru. The species was collected in the Continental rise (563-1,201 m) off Trujillo Department. Reference material has been deposited in the Colección Científica del Instituto del Mar del Perú (IMARPE), Lima, Peru.

**Keywords:** First record, biodiversity, Southeast Pacific.

#### Resumen

Se registra por primera vez para el Perú la especie pelágica de pepino de mar *Enypniastes eximia* Théel, 1882 (Echinodermata: Holothuroidea). La especie fue recolectada en el talud continental (563-1,201 m) frente al departamento de Trujillo. Material de referencia se encuentra depositado en la Colección Científica del Instituto del Mar del Perú (IMARPE), Lima, Perú.

**Palabras clave:** Primer Registro, biodiversidad, Pacífico Sudeste.

Presentado: 26/03/2012  
Aceptado: 28/07/2012  
Publicado online: 01/10/2012

#### Introduction

*Enypniastes eximia* Théel, 1882 (Family Pelagothuridae Ludwig, 1894) is a benthopelagic, cosmopolitan swimming sea cucumber (Miller and Pawson, 1990; Borrero *et al.*, 2003) found at depths between 461 and 5,689 m. This is the first time that *E. eximia* is found in Peruvian waters.

In 1999 Bluhm and Gebruk reported its existence in the "Peruvian Basin" (a circular area of 10.8 km<sup>2</sup>, centered at 7°4.4' S and 88°27.6' W), using photographs of a deep-sea area far away from the Peruvian continental shelf; about 450 nautical miles from the coast (not territorial Peruvian waters).

Specimens were found off Trujillo Department (-7°58.93'S, 80°35.10'W). Organisms were collected using a Bacalao Trawl 586/200 on board of the R/V SHINKAI MARU. Reference material was deposited the Scientific Collection of the Instituto del Mar del Perú (IMARPE).

#### Taxonomy


PHYLUM ECHINODERMATA KLEIN,  
1734 (EX BRUGUIÈRE, 1789)  
ORDER ELASIPODIDA THÉEL, 1882  
SUBORDER PSYCHROPOTINA HANSEN, 1975  
FAMILY PELAGOTHURIDAE LUDWIG, 1894  
GENUS *ENYPNIASTES* THÉEL, 1882

#### *Enypniastes eximia* Théel, 1882

(Figures 1, Table 1)

*Enypniastes eximia*.- Pawson, 1965: 19, 27; Miller & Pawson, 1990: 10-11, figs. 1c,d, 4; Borrero *et al.*, 2003: 76, fig. 3C.

**Material examined:** 450 specimens of three localities (Table 1). IMARPE 03-000475, 1 specimen, August 6<sup>th</sup>, 1999, R/V SHINKAI MARU, Bacalao Trawl 586/200, off Trujillo, between



**Figure 1.** (A) Two specimens of *Enypniastes eximia* Théel, 1882. The large anterior webbed veil is visible in the specimen of the left side. (B) Lateral view. Total length of the specimen ~12 cm.

**Table 1.** *Enypniastes eximia* abundance data (number of specimens and weight). n= number of specimens.

Initial Latitude	07°58.93	08°05.65	06°39.69			
Initial Longitude	80°35.10	80°32.91	81°03.35			
Final Latitude	07°57.52	08°04.03	06°38.53			
Final Longitude	80°35.67	80°33.49	81°04.69			
Depth (m)	1,201	1,163	563			
Date	Aug 6 <sup>th</sup> 1999	Aug 7 <sup>th</sup> 1999	Aug 7 <sup>th</sup> 1999			
Initial time (hrs.)	12:46	16:40	07:45			
Final time (hrs.)	13:16	17:11	08:10			
Trawling time (min.)	30	31	25			
	<i>n</i>	weight (kg)	<i>n</i>	weight (kg)	<i>n</i>	weight (kg)
Total number of specimens and total weight	219	97.56	58	25.78	173	57.65

7°58.93' S, 80°35.10' W and 7°57.52' S, 80°35.67' W, 1,201 m, coll. Y. Hooker; 218 specimens, August 6<sup>th</sup>, 1999, R/V SHINKAI MARU, Bacalao Trawl 586/200, off Trujillo, between 7°58.93' S, 80°35.10' W and 7°57.52' S, 80°35.67' W, 1,201 m, coll. Y. Hooker; 58 specimens, August 7<sup>th</sup>, 1999, R/V SHINKAI MARU, Bacalao Trawl 586/200, off Trujillo, between 8°05.65' S, 80°32.91' W and 8°04.03' S, 80°33.49' W, 1,163 m, coll. Y. Hooker; 173 specimens, August 7<sup>th</sup>, 1999, R/V SHINKAI MARU, Bacalao Trawl 586/200, off Trujillo, between 6°39.69' S, 81°03.35' W and 6°38.53' S, 81°04.69' W, 563 m, coll. Y. Hooker. Capture total weight 180.99 kg.

**Description:** This species has a distinctive barrel-shaped body and a large anterior webbed veil incorporating up to 12 conical podia (Fig. 1A). There are two rectangular postero lateral veils composed of 10-15 webbed podia. No ossicles are found in the body wall or any other part of the external or internal structures. *Enypniastes eximia* is transparent; internal structures, especially the sediment-filled coiled intestine, are readily visible through the body wall. Individuals examined during this study ranged from 11 to 25 cm in length.

**Color:** Reddish to dark purple. The coloration of the integument varies with body size; small individuals are pale pink and large adults are dark brown-red to crimson (Miller and Pawson, 1990).

**Distribution:** *Enypniastes eximia* is cosmopolitan at depths of 461-5,689 m (Miller and Pawson, 1990; Borrero et al., 2003). Swimming specimens observed from submersibles or in seafloor photographs have always been found within a few meters of the seabed (Bluhm and Gebruk, 1999). Billett *et al.* (1985) have collected juvenile specimens as much as 3,000 m above the seabed, and the same species has been reported (as *Pelagothuria bouvieri*) at the surface (Hérouard, 1923).

**Remarks:** *Enypniastes eximia*, employs a “burglar alarm” strategy of predator deterrence. After feeding on benthic sediments it lifts off the bottom and spends most of its time up in the water column. Light production is triggered mechanically and is produced by hundreds of granular bodies within its gelatinous integument. The skin of *E. eximia* is very fragile and strong physical contact causes the skin to be sloughed off in a glowing cloud. The deciduous skin is also sticky and it readily adheres

to most surfaces. Physical contact elicits light production that can “paint” a predator with patches of the glowing skin, thus revealing the presence of the attacker to its own visually cued predators (Robison, 1992).

### Acknowledgments

To the Instituto del Mar del Perú (IMARPE) and Albertina Kameya Kameya (Unidad de Investigaciones en Biodiversidad) to allow us reviewing and work on their scientific collection. To Alicia Duran Gonzalez (ICML, UNAM) for her technical support.

### Literature cited

- Billett D.S.M., B. Hansen & Q.J. Huggett. 1985. Pelagic Holothuroidea of the Northeast Atlantic. In B.F. Keegan and B.D.S. O'Connor (Eds.) Echinodermata: proceedings of the 5th international echinoderm conference galway 24-29 september, 1984, pp. 399-411. Balkema Press, Rotterdam.
- Bluhm H. & A. Gebruk. 1999. Holothuroidea (Echinodermata) of the Peru Basin- Ecological and Taxonomic Remarks based on underwater images. P.S.Z.N. Marine Ecology 20 (2): 167-195.
- Borrero-Pérez G. H., M. Benavides-Serrato, D.O. Solano & G.R. Navas. 2003. Holothuroideos (Echinodermata: Holothuroidea) recolectados en el Talud Continental superior del Caribe Colombiano. Boletín del Instituto Oceanográfico de Venezuela. Univ. de Oriente, 42: 65-85.
- Hansen B. & F. J. Madsen. 1956. On two bathypelagic holothurians from the South China Sea. Scientific Results of the Danish Deep-sea Expedition round the world 1950-52. Galathea Report 2: 55-59.
- Hérouard E. 1923. Holothuries provenant des campagnes des yachts Princesse-Alice et Hirondelle II (1898-1915). Résultats Campagnes Scientifiques Prince-Albert I, 66:1-163.
- Miller J.E. & D.L. Pawson. 1990. Swimming sea cucumbers (Echinodermata: Holothuroidea): A survey, with analysis of swimming behavior in four bathyal species. Smithsonian Contributions to the Marine Sciences, (35): 1-18.
- Pawson D. L. 1965. The bathyal holothurians of the New Zealand Region. Zoology Publications from Victoria University of Wellington (39): 1-33.
- Robison B.H. 1992. Bioluminescence in the benthopelagic holothurian *Enypniastes eximia*. J. Mar. Biol. Assoc. UK 72:463-472.
- Théel, H. 1882. Report on the Holothuroidea, dredged by H.M.S. Challenger during the year 1873-1876, Zoology 4(3): 1-176.